AD_____

REPORT NO T15-89

AD-A209 613

BETWEEN-MEAL FOOD INTAKE FOR RESERVISTS TRAINING IN THE FIELD

U S ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE Natick, Massachusetts

SELECTE JUN 2 1 1969

APRIL 1989



UNITED STATES ARMY
MEDICAL RESEARCH & DEVELOPMENT COMMAND

89 9 20 243

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DISPOSITION INSTRUCTIONS

Destroy this report when no longer needed.

Do not return to the originator.

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188				
1a. REPORT SECURITY CLASSIFICATION				16 RESTRICTIVE MARKINGS				
2a. SECURITY CLASSIFICATION AUTHORITY				3 DISTRIBUTION / AVAILABILITY OF REPORT				
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved funlimited	for public r	elease	e; distribution is		
4. PERFORMIN	IG ORGANIZAT	ION REP	ORT NUMBE	R(S)	5. MONITORING ORGANIZATION REPORT NUMBER(\$)			
6a. NAME OF USARIEM	PERFORMING	ORGANI	ZATION	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION			
				SGRD-UE-NR				Development Cmd
6c. ADDRESS (Natick, M			de)		Fort Detric	ty, State, and ZIP ck MD 21701–50		
8a. NAME OF ORGANIZA		NSORIN	G	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMEN	T INSTRUMENT IC	ENTIFICA	TION NUMBER
8c. ADDRESS (City, State, and	ZIP Cod	le)	<u> </u>	10. SOURCE OF	FUNDING NUMBER	RS	
					PROGRAM ELEMENT NO. 63002D	PROJECT NO. 3M263	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Incl	uda Sacurity C	laccificat	rion)			002D819	A]	DA305222
Between-	Meal Food			eservists Traini	ng in the Fi	ield		
12. PERSONAL M.S. Ros								
13a. TYPE OF Final			13b. TIME CO	OVERED in 88 TO Apr 89	14. DATE OF REPO 1989 April		Day) 1	5. PAGE COUNT 36
16. SUPPLEME	NTARY NOTAT	TION						
17.	COSATI			18. SUBJECT TERMS (
FIELD	GROUP	SUB	-GROUP	Between-Meal F rates. Reservi		reeding, Sr	acks,	MRE consumption
				ruces, Reservi				
Soldiers take non-ration food items to the field to supplement the A-ration and Meals, Ready-to-Eat (MRE) foods that are served. The adequacy of data analyses of nutrient intakes in the field depends on whether the foods eaten between-meals are included in the analyses. During 8 days of work in the field in a hot environment by a reserve field medical unit, the intake of between-meal foods and fluids accounted for 25% of the total energy intake. About 27% of the between-meal foods and fluids were beverages. The soldiers skipped 13% of the A-ration meals with a majority (55.5%) skipping breakfast. MRE consumption for the reservists at the lunch meal in this study (44%) was much lower than the 70-80% reported previously for active duty soldiers when consumption of non-ration between-meal foods was strongly discouraged. 21 ABSTRACT SECURITY CLASSIFICATION WINDLASSIFIED/UNLIMITED SAME AS RPT. DITC USERS Unclassified								
22a. NAME O	RESPONSIBLE	INDIVID		C OTTO OSCRI	226 TELEPHONE	(Include Area Cod		
LTC Madeleine S. Rose			508-651-497	79	SGRI	D-UE-NR		

HUMAN RESEARCH AND DISCLAIMER STATEMENTS

Human subjects participated in these studies after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research.

The views, opinions and findings contained in this report are those of the authors and should not be construed as an official Department of the Army position, policy or decision unless so designed by other official documents.



Acomium For	
VEIS CRASS	A
District 120	ניו
Umarking Jadi	1.1
Jan 1 7 12 1	
27	
Sale to Sale	
and a second	Di Jes
	1143
	11
A	
A-1	

BETWEEN-MEAL FOOD INTAKE FOR RESERVISTS TRAINING IN THE FIELD

LTC Madeleine S. Rose

U.S. Army Research Institute of Environmental Medicine
Natick, MA 01760-5007

April 1989

FOREWORD

The information contained in this technical report is derived from the data collected for the technical report, Effectiveness and Acceptability of Nutrient Solutions in Enhancing Fluid Intake in the Heat. Complete food and fluid intake were collected for soldiers working for 8 days in the heat. The data were collected in sufficient detail to make it possible to analyze the data for A-ration meals, Meal, Ready-to-Eat (MRE) rations, and between-meal foods.

TABLE OF CONTENTS

Foreword	iii
Table of Contents	iv
List of Table	v
Abstract	vi
Introduction	1
Methods	3
Results and Discussion	5
Conclusion	11
References	12
Appendix A - MREs Eaten during the Field Exercise	14
Appendix B - Frequency Distribution by Category of Foods Eaten Between-Meals in the Field	15
Appendix C - Between-Meal Foods Eaten during the AM Period	16
Appendix D - Between-Meal Foods Eaten during the PM Period	20

LIST OF TABLES

NUI	MBER	PAGES
1	Physical characteristics of subjects	5
2	Distribution of subjects according to the number of meals skipped	6
3	Contribution to total energy intake of A-ration meals, MRE rations, and between-meal foods	9

INTRODUCTION

The food intake of soldiers in the field has been studied repeatedly (1-8). However, the completeness of the data is sometimes questionable. Anecdotal stories indicate that soldiers do not restrict their food intake to the rations that are served in the field, but bring their own favorite foods to supplement the rations. Complete food consumption data usually are collected on the military rations; however information on the extra foods eaten between meals is more difficult to collect and quantitate. This source of nutrient intake was not collected and/or analyzed in some field (4,9) and dining facility studies (10-12). Some field studies have controlled the sources of between-meal food intakes by checking all rucksacks for unauthorized food (7) or by utilizing the chain of command to strongly discourage the consumption of non-issued food items in the field (1). If between-meal food data are collected, their contributions to overall nutrient intakes are included in the totals. The study by Roberts et al. (6) was one of the few ration studies that reported the nutrient intakes from non-ration food items separately. Published data are generally not available on the types and the amounts of between-meal foods that are eaten in the field. The objective of this technical report was to describe the types and amounts of between-meal foods that

were eaten by reserve soldiers working 8 days in the field in a hot environment.

METHODS

The data for this technical report were collected as part of a larger study on the acceptability and effectiveness of the carbohydrate-electrolyte solutions in enhancing fluid intake in the heat (3). The methods are reported in detail by Rose et al. (3); however, the methods are described briefly here. Volunteers were recruited from reservists of the 44th Evacuation Hospital, 807th Medical Brigade, participating in a field training exercise at Fort Hood, TX during June 1988. Data from 61 subjects are included in this data analysis even though 6 of the 61 subjects did not complete the entire study. In general, the activity level of the soldiers averaged out to moderate activity. There were spurts of heavy and light activity as the soldiers erected and lived in tents in the field for the 8 days of the exercise. Some of the soldiers attended classes in garrison during the day. On Day 5 of the study, all soldiers, including the test subjects, were allowed to return to garrison in the afternoon to shop at the PX, take showers, attend movies, etc. The subjects returned to the field by 2200 hours that night.

The subjects were allowed to consume foods and fluids <u>ad</u>

<u>libitum</u>. Data were collected for the: a) A ration meals served

at the breakfast and dinner meals (dinner changed to lunch on day

5); b) Meals, Ready-to-Eat version 6 (MRE) for lunch; c) between
meal foods and fluids eaten during the period 1630-0530 hours (PM

card); and d) between-meal foods and fluids eaten during the period 0530-1630 hours (AM card) except Day 5 when the data from the AM and PM cards were combined and collected at 0530 hours on Day 6. Information on food and fluid intake at the two hot Aration meals was collected by the nutrition data collectors using a modified visual estimation method (13) developed at the U.S. Army Research Institute of Environmental Medicine (USARIEM). this method, the subject showed his food selections to the data collector, ate his meal, and showed his tray to the data collector again. The data collector compared the food portions to pre-weighed standards to estimate portion size. Information on the intake of the MRE were recorded separately by the subjects on the front of the food and fluid intake cards. All the MRE food components were listed on the food and fluid intake cards and the subjects were asked to circle the food and amount eaten. The MREs were issued to the soldiers at the breakfast meal to be eaten at the lunch meal. On Day 5 the breakfast and lunch meals were A-rations and the MRE was issued for the dinner meal. All extra foods brought to the field, purchased from the PX mobile kitchen, and eaten at restaurants or fast food establishments were recorded on the back of the between-meal food and fluid intake cards. Data were collected on 100% of the A-ration meals served in the field and only 4% of the between-meal data forms were missing. The missing data will cause the energy and fluid intake to be slightly underestimated.

RESULTS AND DISCUSSION

Physical characteristics of subjects participating in this study are listed in Table 1. The weights in the table are the weights of the reservists at their home station (Oklahoma City, OK or El Paso, TX) before the soldiers were transported to Fort Hood, TX. Some data were missing on height because some soldiers did not fill in the final questionnaire with the demographic data. The subjects lost less than 1 kg body weight over 8 days and therefore were very close to ingesting their energy requirements. The subjects were almost evenly divided by gender and military rank. About 45% of the volunteers had served in the military for less than 5 years.

Table 1. Physical characteristics of subjects.

VARIABLE	NUMBER	MEAN+SEM	MINIMUM	MUMIXAM
Age, years	61	33.6 <u>+</u> 1.1	19	51
Height, cm	51	173.5 + 1.5	152	198
Weight, kg*	61	75.4 <u>+</u> 2.0	41.5	128.2

^{*}Pre-deployment weight

As reported previously (3), 13% of the field meals were skipped deliberately (Table 2) and most were breakfast meals.

About 55.5% of the skipped meals were breakfast meals and 44.5% were dinner meals. Since all soldiers were wakened for formation just prior to the breakfast meal most of the skipped breakfasts

were probably deliberate. The hot A-ration meals were popular since 19 of the 61 subjects ate all the A-ration meals that were served in the field. Most of the soldiers (85%) missed 3 meals or less of a possible 16 (2 meals/day x 8 days) meals. Data show that one person skipped 11 meals and 1 person skipped 15 meals (Table 2). The subject who missed 15 meals had come prepared to skip all meals. She was attempting to lose weight and had brought her own food: yogurt, distilled water, skim milk, etc.

Table 2. Distribution of subjects according to the number of meals skipped.

NUMBER	OF	MEALS	SKIPPED	ħ	NUMBER	OF	SUBJECTS
		0				19	
		1				13	
		2 3				10	
						10	
		4				2	
		5				2	
		6 7					
		8				1	
		9				(
		10				(
		11				1	
		12				ď	
		13				Č	
		14				Č	
		15				ì	_
		16				Ċ)
	7	TOTAL				61	_

Adapted from reference #3.

The subject who missed 11 meals was coordinating administrative details of the FTX and so skipped many of the meals served in the field, but he was recording his food intake on the between-meal food and fluid intake card.

The mean intake for the 26 males who completed all 8 days of the study (3) was 3056 ± 74 kcal/day which is less than the Military Recommended Dietary Allowances (MRDA) of 3200 kcal/day set for moderately active male military personnel, ages 17 to 50 years (14). Compared to the mean male intakes, the intakes of the 29 females were much lower at 2343±55 kcal/day. The daily caloric intakes ranged from a minimum of 137 kcal/day for one subject to a maximum of 6162 kcal/day for another subject. The mean male energy intake for the present study is within the range the energy intake of previous field studies where mean intakes of subjects consuming 2 A-ration meals + 1 MRE ration/day for extended periods (5-6 weeks) have been reported to be 2950 and 3271 kcal/day (1,15). A mean intake of 3713 kcal/day was reported for field artillery soldiers fed 3 A-ration meals during 8 days of sustained artillery operations in the field (2). subjects in the present study were very close to equilibrium in their energy intake and expenditure since the mean weight loss was less than 1 kg. This group of subjects may not have needed excessive calories because they started with moderate activity (with spurts of heavy activity) during the first 3 to 4 days while setting up the hospital area, but their activity level

decreased markedly during the later days while they waited for events to occur in the FTX.

The frequency of MRE consumption is shown in Appendix A.

Only about 44% of the MREs which should have been consumed in the field were eaten (215 MRE entrees consumed/8 days x 61 subjects). This percentage was very low considering that the CFFS-FDTE (1) study had shown that male and female soldiers ate about 80 and 70% of the MREs, respectively. This apparent unwillingness to eat MREs could account for some of the inadequate nutrient intakes found in this study since one MRE supplies 1200 kcal and ~1/3 of the MRDA for one meal. However, the tables (Appendix B) showing the frequency of between-meal food consumption indicate that the soldiers were not suffering from a caloric deficit.

The tables in Appendix B show the frequency of between-meal food consumption in the field. Between-meal foods include: foods brought to the field site by the reservists and foods purchased in restaurants, PX, fast food establishments, etc.

Some of this between-meal food was provided by the PX mobile kitchen which made regular visits at the field site. Other foods were eaten while the soldiers were on-post on Day 5 or other days for classes. The energy values in Table 3 of the present report are slightly higher than those reported on the same subjects by Rose et al. (3) because the data in the previous technical report are based on only 55 subjects that completed all 8 days of the study. The contributions of the different food sources (A-

ration, MRE, between-meal) to overall energy intake was similar for males and females. Approximately 50% of the total energy intake was contributed by the A-ration meals, 25% from the MREs, and another 25% from the between-meal foods (Table 3). Since the energy intakes of the subjects in the present study were approximately the same as those of previous field studies (1,2,15), the caloric contribution from between-meal foods indicates that the soldiers reduced their intake of A-ration and MRE calories by 25%.

Table 3. Contribution to total energy intake of A-ration meals, MRE rations, and between-meal foods.

FOOD SOURCE	TOTAL	MALES	FEMALES
	KCAL (%)	KCAL (%)	KCAL (%)
A-RATION MEALS	1528±30 (49%) ^a 804±27 (26%) ^b 761±30 (25%) ^b	1645 <u>+</u> 43 (49%)	1404 <u>+</u> 41 (51%)
MRE RATION		892 <u>+</u> 36 (26%)	672 <u>+</u> 38 (24%)
BETWEEN-MEAL FOOD		846 <u>+</u> 50 (25%)	682 <u>+</u> 35 (25%)
TOTAL (n=61)	3093	3383	2758

Values are Mean±SEM Means with unlike superscripts differ, p<0.001.

Anecdotal reports have indicated that all soldiers took foods to the field to supplement MRE meals, but no other study has generated data on the types of between-meal foods (Appendices C and D) and their contribution to the overall energy intake of soldiers working for extended periods of time in the field. The

Reservists were not restricted in the types of foods that they could take to the field because the investigators wanted normal food consumption during the field exercise. Between-meal food and fluid intakes averaged about 3.74 snacks/day/person. It is interesting that about 27% of all between-meal foods eaten in the field were fluids. A reason for the high beverage consumption could be an attempt to increase fluid intake in the heat. soldiers participating in the present study were part of a medical unit that had received extensive training on water discipline and the importance of preventing dehydration during work in the heat. The consumption of foods from the fruit/juice, meat, snack/chips, candy/gum, bread/cereal, and dessert categories ranged from 8.1-14.6%. Less than 100 servings of the 1823 between-meal food items were consumed in each of the other categories. Most of the between-meal food items were canned or dried and did not require any refrigeration. However, a few items (yogurt, dairy products, skim milk) did require cold storage.

The results show that between-meal food intake should be included in all analyses of nutrient intake. Assuming that the amount of food that soldiers take to the field is minimal would underestimate nutrient intakes. The 25% contribution from between-meal foods in this study is probably higher than would be typical for active duty troops due to the nature of the unit studied (access to between-meal foods because of proximity to

transportation, classes in garrison, PX mobile kitchen, etc.); however, it does show that non-ration foods can affect total energy intake for soldiers eating A-rations and MREs in the field.

CONCLUSIONS

A major percentage of the total energy intake came from non-ration food items. About 25% of the total energy consumed in the field was from between-meal foods/fluids and another 25% from MREs. Without these between-meal foods, energy intake would have been inadequate, but it cannot be determined whether having the extra between-meals foods caused the reduction of A-ration and MRE food intake at meals or whether having the between-meal foods increased inadequate intakes to adequate levels for the soldiers in the present study. The soldiers skipped 13% of the A-ration meals with the majority (55.5%) of the skipped meals at breakfast. MRE consumption for the reservists at the lunch meal in this study (44%) was much lower than the 70-80% reported previously for active duty soldiers when consumption of non-ration between-meal foods was strongly discouraged.

REFERENCES

- 1. US Army Combat Developments Experimentation Center, Fort Ord, CA and US Army Research Institute of Environmental Medicine, Natick, MA. Combat Field Feeding System-Force Development Test and Experimentation (CFFS-FDTE) Test Report CDEC-TR-85-006A, 1986.
- 2. Rose MS, Carlson DE. Effects of A-ration meals on body weight during sustained field operations. (Technical Report No. T2-87) Natick, MA: US Army Research Institute of Environmental Medicine, 1986.
- 3. Rose MS, Szlyk PC, Francesconi RP, Lester LS, Armstrong L, Matthew W., Cardello AV, Popper RD, Sils I, Thomas G, Schilling D, Whang R. Effectiveness and Acceptability of nutrient solutions in enhancing fluid intake in the heat. (Technical Report No. T10-89) Natick, MA: US Army Research Institute of Environmental Medicine, 1989.
- 4. Edinberg J, Engell D. Field evaluation of the B ration in a hot weather environment. (Technical Report No. TR-89/002) Natick, MA: US Army Natick Research, Development and Engineering Center, 1988.
- 5. Morgan TE, Hodgess LA, Schilling D, Hoyt RW, Iwanyk EJ, McAninch G, Wells TC, Hubbard VS, Askew EW. A comparison of the meal, ready-to-eat, ration, cold weather, and ration, lightweight nutrient intakes during moderate altitude cold weather field training operations. (Technical Report No. T5-89) Natick, MA: US Army Research Institute of Environmental Medicine, 1988.
- 6. Roberts DE, McGuire BJ, Engell DB, Salter CA, Rose MS. The role of water consumption on consumption of the ration, cold weather. (Technical Report In Preparation) Natick, MA: US Army Research Institute of Environmental Medicine.
- 7. Askew EW, Munro I, Sharp MA, Siegel S, Popper R, Rose MS, Hoyt RW, Martin JW, Reynolds K, Lieberman HR, Engell D, Shaw CP. Nutritional status and physical and mental performance of special operations soldiers consuming the ration, lightweight or the Meal, ready-to-eat military field ration during a 30-day field training exercise. (Technical Report No. T7-87) Natick, MA: US Army Research Institute of Environmental Medicine, 1987.

- 8. Roberts DE, Askew EW, Rose MS, Sharp MA, Bruttig S, Buchbinder JC, Engell DB. Nutritional and hydration status of special forces soldiers consuming the ration, cold weather or the meal, ready-to-eat ration during a ten day cold weather field training exercise. (Technical Report No. T8-87) Natick, MA: US Army Research Institute of Environmental Medicine, 1987.
- 9. Glassford DL, Ponsioen AM, Prevo C, Szeto EG. Concept evaluation program (CEP) test of the combat field feeding system, (medical). US Army Medical Dept Board, AHS, Fort Sam Houston, TX. Project No. 4-86, May 1987.
- 10. Szeto EG, Carlson DE, Dugan TB, Buchbinder JC. A comparison of nutrient intakes between a Ft. Riley contractor-operated and a Ft. Lewis military-operated garrison dining facility. (Technical Report No. T2-88) Natick, MA: US Army Research Institute of Environmental Medicine, 1987.
- 11. Szeto EG, Dugan TB, Gallo JA. Assessment of habitual diners nutrient intakes in a military-operated garrison dining facility Ft. Devens I. (Technical Report No. T3-89) Natick, MA: US Army Research Institute of Environmental Medicine, 1988.
- 12. Szeto EG, Gallo JA, Samonds KW. Passive nutrition intervention in a military-operated garrison dining facility FT Devens II. (Technical Report No. T7-89) Natick, MA: US Army Research Institute of Environmental Medicine, 1989.
- 13. Rose MS, Buchbinder JC, Dugan TB, Szeto EG, Allegretto JD, Rose RW, Carlson DE, Samonds KW, Schnakenberg DD. Determination of nutritional intakes by a modified visual estimation method and computerized nutrition analysis for dietary assessments of military field and garrison feeding. (Technical Report No. T6-88) Natick, MA: US Army Research Institute of Environmental Medicine, 1987.
- 14. Army Regulation 40-25. Nutrition Allowances, Standards, and Education. Headquarters, Departments of the Army, the Navy, and the Air Force. Washington, DC, 15 May 1985.
- 15. Hirsch E, Meiselman HL, Popper RD, Smith G, Jezior B, Lichton I, Wenkam N, Burt J, Fox M, McNutt S, Thiele MN, Dirige O. The effects of prolonged feeding meal, ready-to-eat (MRE) operational rations. USANRDC Technical Report No. TR-85/035, 1983.

APPENDIX A MRES EATEN DURING THE FIELD EXERCISE

FOOD ITEM NAME	CODE	FREQUENCY
APPLESAUCE, MRE BEEF PATTIES, MRE BEEF W/BBQ SAUCE, MRE BEEF STEW, MRE BEEF W/GRAVY, MRE BEEF W/SPICED SAUCE, MRE BEANS W/TOMATO SAUCE, MRE CANDY (ALL TYPES), MRE CATSUP, MRE CHICKEN A LA KING, MRE	MR6601	14
REEF PATTIES, MRE	MR6303	15
BEEF W/BBO SAUCE, MRE	MR6304	36
BEEF STEW, MRE	MR6305	2
BEEF W/GRAVY, MRE	MR6308	27
BEEF W/SPICED SAUCE, MRE	MR6312	3
BEANS W/TOMATO SAUCE, MRE	MR6313	24
CANDY (ALL TYPES), MRE	MR6815	30
CATSUP, MRE	MR6904	5
CHICKEN A LA KING, MRE	MR6309	30
CHOC CVD COOKIE BAR, MRE	MR6801	65
CHOC CVD BROWNIE, MR	MR6803	40
CHERRY NUT CAKE, MRE	MR6804	4
CHOC NUT CAKE, MRE	MR6807	2
CHEESE SPREAD, MRE	MR6901	53
COCOA POWDER, MRE	MR6001	19
COFFEE, MRE	MR6002	10
CREAM SUBSTITUTE, MRE	MR6003	7
CRACKERS, MRE	MR6701	200
FRANKFURTERS, MRE	MR6306	19
FRUIT MIX, MRE	MR6604	46
FRUITCAKE, MRE	MR6806	30
GRAVY BASE (SOUP MIX), MRE	MR6905	1
GUM, MRE	MR6816	58
HAM/CHICKEN LOAF, MRE	MR6302	13
JELLY, MRE	MR6903	45
MAPLE NUT CAKE, MRE	MR6805	23
MEATBALLS W/BBQ SAUCE, MRE	MR6310	28
PEACHES, MRE	MR6603	46
PEANUT BUTTER, MRE	MR6906	63
PEARS, MRE	MRI609	12
PORK SAUSAGE PATTIES, MRE	MR6301	15
POTATO PATTY, MRE	MR6401	1
SALT, MRE	MR6910	15
STRAWBERRIES, MRE	MR6602	1
SUGAR, MRE	MR6909	6
CATSUP, MRE CHICKEN A LA KING, MRE CHOC CVD COOKIE BAR, MRE CHOC CVD BROWNIE, MR CHERRY NUT CAKE, MRE CHOC NUT CAKE, MRE CHOC NUT CAKE, MRE CHEESE SPREAD, MRE COCOA POWDER, MRE COFFEE, MRE CREAM SUBSTITUTE, MRE CRACKERS, MRE FRANKFURTERS, MRE FRUIT MIX, MRE FRUIT MIX, MRE GRAVY BASE (SOUP MIX), MRE GUM, MRE HAM/CHICKEN LOAF, MRE JELLY, MRE MAPLE NUT CAKE, MRE MEATBALLS W/BBQ SAUCE, MRE PEACHES, MRE PEANUT BUTTER, MRE PEARS, MRE PORK SAUSAGE PATTIES, MRE POTATO PATTY, MRE SALT, MRE STRAWBERRIES, MRE SUGAR, MRE TURKEY W/GRAVY, MRE	MR6307	27

APPENDIX B FREQUENCY DISTRIBUTION BY CATEGORY OF FOODS EATEN BETWEEN-MEALS IN THE FIELD

CATEGORY		
FREQUENCY	PERCENT	
Beverages	528	29.0
Fruit/Juices	267	14.6
Meat	160	8.8
Snacks/Popcorn/Chips	159	8.7
Candy/Gum	149	8.2
Breads/Crax/Rice Cakes/Cereal	148	8.1
Desserts/Cakes/Cookies/Puddings	106	5.8
Seeds/Nuts	79	4.3
Dairy Products/Cheese/Yogurt	73	4.0
Fat/Cream Cheese/Salad Dressing	29	1.6
Peanut Butter/Cheese Crax	26	1.4
Pastry	24	1.3
Condiments	23	1.3
Granola	23	1.3
Hamburger/Cheeseburger	13	0.7
Vegetables	13	0.7
Soup	3	0.2

APPENDIX C BETWEEN-MEAL FOODS EATEN DURING THE AM PERIOD (0530-1630 HR)

FOOD ITEM	FREQUENCY
APPLE, RAW APPLESAUCE	61 5
BACON, OVEN FRIED BANANA, FRESH BEEF JERKEY BEEF STICK BISCUITS REFRIGERATED BOLOGNA BREAD, ASST BUBBLE GUM, SUPER	3 7 16 2 1 2 18 3
CAKE W/VAN ICING CANDY, ASST CARROT, RAW, STICKS CATFISH, BREADED CATSUP CELERY, RAW, SLICED CEREAL, ASST CHEESE SPREAD CHEESE BURGER/HAMBURGER CHEESE DANISH CHEESE DANISH CHEX SNACK MIX CHICKEN, CANNED CHICKEN, OVEN FRIED CHICKEN FRIED STEAK CHICKEN MCNUGGETS, MCDONALD CHICKEN PLANKS CHICKEN SALAD CHILI, VENDING CHIPPED BEEF/DRIED CHIPS, ASST SNACK CHOCOLATE CHIPS COFFEE, BREWED	2 71 1 1 5 1 36 4 9 1 1 2 3 3 1 1 1 1 2 42 1 32
COFFEE CREAMER COOKIE, ASST CORN NUT SNACK COTTAGE CHEESE COUGH DROPS CRACKERS, ASST CREAM CHEESE CUPCAKE, CREAM FILLED CUP-A-SOUP, BEEF VEG	1 52 4 2 1 52 6 1

BETWEEN-MEAL FOODS EATEN DURING THE AM PERIOD (0530-1630 HR) (cont'd)

FOOD ITEM	FREQUENCY
DEVILED HAM DINNER ROLL DONUT, FILLED, PLAIN DRESSING, ASST, REG	5 1 4 10
EGG, BOILED or SCRAMBLED	4
FRUIT COCKTAIL IN JUICE FRENCH FRIES FRUIT ROLL/ROLLUP FRUIT, MIXED AND DRIED FUNYON'S ONION RINGS	5 7 3 7 1
GATORADE GRANOLA BAR, ASST GRAPEFRUIT, FRESH GREEN CHILI SAUCE GUM, REGULAR	1 12 1 1 3
HAM, SLICED HAM SANDWICH W/CHEES HERRING, SMOKED, KIPPERS HOAGIE, SANDWICH HONEY HOT DOG, FOOT LONG HOT DOG, FOOT LONG, CHILI HOT SAUCE HUSH PUPPIES	1 3 3 4 1 8 6 1
ICE CREAM,	4
JELLY, ASSORTED FLAV JUICE, ASST	4 20
KOOLAID, ANY FLAVOR	15
LETTUCE	6
MARGARINE, PATTIE MCDONALD'S SAUSAGE & EGG MCMUFFIN MELBA TOAST, 1 PIECE MILK, CHOCOLATE, UHT MILK, WHITE, UHT 2% MILK, SKIM, PROTEIN	1 4 9 18 3

BETWEEN-MEAL FOODS EATEN DURING THE AM PERIOD (0530-1630 HR) (cont'd)

FOOD ITEM	FREQUENCY
MUFFIN, BLUEBERRY	1
NUTS, ASST	42
ONION RINGS ORANGE, WHOLE	1 60
PEACHES, HEAVY SYRUP PEANUT BUTTER SANDWICH PEANUT BUTTER PEARS, CANNED, IN WATER PEPPER, JALAPENO PIE, FRIED, FRUIT FLAVORED PINEAPPLE CHUNKS, CANNED PICO DIGALLO PIZZA, ASST PLUM, FRESH POPCORN, BUTTERED & SALTED POPCORN, SUGAR COATED POP TARTS, FROSTED PORK RINDS/SKINS POTATO SALAD PORK & BEANS, CANNED PRUNES PUDDING, SNACK PACK	6 2 16 1 2 3 1 18 1 14 6 3 2 1 7 5 4
RAISINS RAISIN, YOGURT COATED RAISIN BREAD RICE RICE CAKES, UNSALTED ROLL, CINNAMON ROAST BEEF, COOKED	11 1 1 2 2 2
SALT, INDIV PACKET SARDINES, IN OIL SAUSAGE, SUMMER SAUSAGE SAUSAGE, VIENNA SODA, ASST SODA, DIET SOUP, READY MADE STEAK SAUCE SUNFLOWER SEED KERNEL SUNKIST FUN FRUITS	4 2 1 5 13 180 85 2 1 5

BETWEEN-MEAL FOODS EATEN DURING THE AM PERIOD (0530-1630 HR) (cont'd)

FOOD ITEM	FREQUENCY
SUGAR, GRANULATED SWISS COFFEE DRINK	3 1
SWEET AND LOW	ī
TACO SAUCE	2
TACO SALAD TEA, FROM BAG	1 4
TEA, ICED, PRESWEETENED TORTILLA, FLOUR, 6IN	6 3
TOMATO, RAW, SLICED TRAIL MIX	2 10
TUNA IN OIL	1
TUNA SALAD SANDWICH TUNA IN WATER	1 14
TURKEY, CANNED	1
YOGURT, WEIGHT WATCHERS	4 2
YOGURT, ASSORTED FRUIT	_
ZINGERS, RASPBERRY	2

APPENDIX D BETWEEN-MEAL FOODS EATEN DURING THE PM PERIOD (1630-0530 HR)

FOOD ITEM	FREQUENCY
APPLE APPLESAUCE	22 1
BACON BITS BANANA, FRESH BEEF STICK BEEF JERKEY BIG MAC, MCDONALD'S BREAD, WHITE, SOFT BUBBLE GUM, SUPER	1 2 1 9 2 5 1
CAKE W/CHOC ICING CANDY, ASST CANTALOUPE, DICED CARROT, RAW, STICKS CEREAL, ASST CHEESE SPREAD CHEESE SPREAD CHEESEBURGER CHEESE, CHEDDAR CHEESE CURLS/CHEETOS CHEESE SANDWICH CRACKERS CHEX SNACK MIX CHICKEN, CANNED CHICKEN, CANNED CHICKEN FRIED STEAK CHIP DIP, ASST CHIPS, ASST CINNAMON NUTROLL COFFEE, BREWED COOKIE, ASST CORN NUT SNACK CRACKERS, ASST	1 47 2 1 5 1 2 2 10 1 3 2 1 6 24 1 8 3 8 6 6
DEVILED HAM DONUT, CINNAMON	1 3
FILET MIGNON FRENCH FRIES FRUIT COCKTAIL IN HEAVY SYRUP FRUIT COCKTAIL IN JUICE FRUIT, DRIED FRUIT ROLL/ROLLUP FUNYON'S ONION RINGS	1 2 1 1 1 2

BETWEEN-MEAL FOODS EATEN DURING THE PM PERIOD (1630-0530 HR) (cont'd)

FOOD ITEM	FREQUENCY
GATORADE GRAVY, CREAM GRANOLA BAR , ASST GREEN CHILI SAUCE	2 1 11 1
HAMBURGER HAM, SLICED HERBAL TEA, ANY FLAVOR HOT DOG, FOOT LONG	1 1 1 3
ICE CREAM BAR, CHOCOLATE ICED TEA	1 2
JELLY, ASSORTED FLAVORS JUICE, CRAN-RASPBERRY COCKTAIL JUICE, CRANTASTIC JUICE, GRAPEFRUIT JUICE, ORANGE	2 1 1 3 5
KOOLAID, UNSWEETENED KOOLAID, ANY FLAVOR	1 16
MARGARINE, PAT MILK, WHOLE MILK, CHOCOLATE, UHT MILK, WHITE, UHT 2%	2 2 8 9
NUTS, ASST NUTS, CASHEWS NUTS, PEANUTS, HONEY ROASTED	19 5 5
ORANGE	19
PEANUT BUTTER SANDWICH PEANUT BUTTER PEACH, FRESH PEARS, CANNED PIE, FRIED, FRUIT FLAVORED PIZZA, PEPPERONI POPCORN, BUTTERED & SALTED POPCORN, CHEESE FLAVORED POPCORN, SUGAR COATED POP TARTS, FROSTED, ASST PORK RINDS/SKINS POUND CAKE	4 3 1 1 2 1 21 1 3 3 1 1

BETWEEN-MEAL FOODS EATEN DURING THE PM PERIOD (1630-0530 HR) (cont'd)

FOOD ITEM	FREQUENCY
PRETZELS PRUNES PUDDING, SNACK PACK	6 1 2
RAISIN RICE CAKES, UNSALTED	1
SARDINES, IN OIL SAUSAGE, SUMMER SAUSAGE, VIENNA SNACK CAKE SODA, ASST SODA, DIET SOUR CREAM STRAWBERRIES, FRESH SUNFLOWER SEED KERNELS SWISS COFFEE DRINK SWEET AND LOW	1 2 3 2 115 50 2 1 8 1
TEA, FROM BAG TEA, ICED, PRESWEETENED TRAIL MIX TUNA SALAD FOR SANDWICH	1 7 6 1
V-8 JUICE	3
WEIGHT WATCHER'S WAFERS	3
YOGURT, ASSORTED FRUIT YOGURT, WEIGHT WATCHERS	1 2

DISTRIBUTION LIST

	NO.	OF	COPIES
Defense Technical Information Center ATTN: DTIC-DDA Alexandria, VA 22304-6145			19
Commander U.S. Army Medical Research and Development Command SGRD-RMS SGRD-PLC Fort Detrick Fredrick, MD 21701-5012			1 1
Commandant Academy of Health Sciences, U.S. Army ATTN: AHS-CDM ATTN: HSHA-CDM ATTN: HSHA-CDS Fort Sam Houston, TX 78234			1 1 1
Dir of Biol & Med Sciences Division Office of Naval Research 800 N. Quincy Street Arlington, VA 22217			1
CO, Naval Medical R&D Command National Naval Medical Center Bethesda, MD 20014			1
HQ AFMSC/SGPA Brooks AFB, TX 78235			1
Under Secretary of Defense Research and Engineering ATTN: OUSDRE(RAT)E&LS Washington, DC 20310			1
Dean School of Medicine Uniformed Services University of Health Sciences 4301 Jones Bridge Road Bethesda, MD 20014			1
Commander U.S. Army War College Carlisle Barracks, PA 17013			1
Commander U.S. Army Soldier Support Center Ft. Benjamin Harrison, IN 46216			1

DISTRIBUTION LIST (continued)

NO. O	e COPIES
Assistant Secretary of Defense (Health Affairs) ATTN: ASD(HA) PA&QA Washington, DC 20310	1
Assistant Secretary of Defense (Aquisition & Logistics) ATTN: OASD(A&L)SD Washington, DC 20310	1
Commander U.S. Army Troop Support Command ATTN: AMSTR-E 4300 Goodfellow Boulevard St. Louis, MO 63120-1798	1
Commander U.S. Army Test and Evaluation Command ATTN: AMSTE-EV-S Aberdeen Proving Ground, MD 21005-5055	1
Commander U.S. Army Operational Test Evaluation Agency ATTN: CSTE-ZX 5600 Columbia Pike Falls Church, VA 22041	1
Commander U.S. Army Training and Doctrine Command ATTN: ATCD-S Fort Monroe, VA 23651	1
Commander U.S. Army TRADOC Combined Arms Test Activity ATTN: ATCT-PO Ft. Hood, TX 76544	1
Commander U.S. Army Materiel Command ATTN: AMCDE-S Alexandria, VA 22333	1
Commander U.S. Army Combined Arms Center ATTN: ATZL-TIE Fort Leavenworth, KS 66027-5130	1

DISTRIBUTION LIST (continued)

	NO.	OF	COPIES
HQDA OTSG ATTN: DASG-DBD Rm 617, Bldg 5 Skyline Place 5111 Leesburg Pike Falls Church VA 22041-3258			1
HQDA ATTN: DASG-RDZ Washington, DC 20310-2300			1
HQDA DCSLOG ATTN: DALO-TST Washington, DC 20310-2300			1
Commandant U.S. Army Quartermaster School ATTN: ATSM-CDT ATTN: ATSM-SFS-FM Fort Lee, VA 23807			1
Commandant U.S. Army Troop Support Agency ATTN: DALO-TAF ATTN: DALO-TAF-F FT. Lee, VA 23801			1
Commander U.S. Army Natick Research, Development and Engineering Center ATTN: STRNC-W ATTN: STRNC-Y ATTN: STRNC-T ATTN: STRNC-E ATTN: STRNC-E ATTN: STRNC-E ATTN: STRNC-TAA Natick, MA 01760-5000			1 1 1 1
HQ U.S. Marine Corps Code LFS-4 Washington, DC 20380-0001			1

DISTRIBUTION LIST (continued)

	NO.	OF	COPIE
Dept of Clinical Investigation Chief, Army Medical Specialist Corp-CIS WRAMC			2
Washington, DC 20307-5001			
Commander U.S. Army Training and Doctrine Command ATTN: ATPL-MSS Fort Monroe, VA 23651-5000			1
MAJ Robert Stretch DCIEM 1133 Sheppard Ave. West P.O. Box 2000 Downsview, Ontario, Canada M3M 3B9			2
HQ, V Corps ACofS, G1 ATTN: AETV-GAD, Lynne Man, RD, MPH APO NY 09079			2
Health/Fitness Nutritionist ATTN: Dr. Bernadette Feist-Fite NDH-A-ED Fort McNair, DC 20319-6000			1
U.S. Army Physical Fitness School ATTN: Ms. Terrie Clarke, RD Fort Benjamin Harrison, IN			1
Dietitian Staff and Faculty - Cadet Mess US Military Academy West Point, NY 10996			1

UNOFFICIAL MAILING LIST

H.E. Sauberlich, Ph.D.
Professor and Director
Division of Experimental Nutrition
Dept. of Nutrition Sciences
University of Alabama
Birmingham, AL 35294

J.E. Canham, M.D. Director, Medical Affairs Kabi Vitrum 1311 Harbor Bay Parkway Alameda, CA 94501

Current Members Committee on Military Nutrition Selected Past Members:

Mark Hegsted, Ph.D. Professor Nutrition Emeritus 58 Boulder Road Wellesley Hills, MA 02181

A.L. Hecker, Ph.D.
Director Medical Nutrition Research
Department of Medicine (480)
Ross Laboratories
625 Cleveland Ave
Columbus, Ohio 43215

J.M. Iacono, Ph.D.
Director Western Human Nutrition Res. Ctr.
Agr. Res. Serv.
USDA
P.O. Box 29997
Presidio of San Francisco, CA 94129

Patricia Deuster, Ph.D. Dept. of Mil. Med. U.S.U.H.S. 4301 Jones Bridge Road Bethesda, MD 20814-4799

M.J. Kretsch, Ph.D.
Western Human Nutrition Res. Ctr.
Agr. Res. Serv.
USDA
P.O. Box 29997
Presidio of San Francisco, CA 94129

MAJ Judith Turcotte SGPS-FP 5109 Leesburg Pike Rm 606 Falls Church, VA 22041-3258

LTC Craig Palmer Nutrition Care Branch (HSHA-IHC) AHS Fort Sam Houston, TX 78234-6200

Dr. Gerald F. Combs, Ph.D.
Asst. Deputy Administrator, Human Nutrition
ARS, USDA
Rm 132, BARC-W
Beltsville, MD 20705

Stimson Library
Academy of Health Sciences
ATTN: Ms. Kay Livingston
Bldg. 2840, Rm 106
Fort Sam
Houston, TX 78234~6100